



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

I. *A Catalogue of the Fifty Plants from Chelsea-Garden, presented to the Royal Society, by the Company of Apothecaries, for the Year 1725, pursuant to the Direction of Sir Hans Sloane, Baronet, Pr. Coll. Med. S. P. V., Pr. By Mr. Isaac Rand, Apothecary, F. R. S.*

- 151 **A** Lchimilla vulgaris, C.B. 319.  
 152 — Alpina, pubescens, minor, H. Reg. Par.  
 153 After Americanus Belvidere foliis, floribus ex cœruleo albicantibus, spicis prælongis. Pluk. Phyt. Tab. 78. f. 5.  
 154 — Virginianus, angustifolius, ferotinus, parvo albente flore, Park. Th. 132.  
 155 — —, pyramidatus, Hyssopi foliis, asperis, calycis squamulis foliaceis.  
 156 — maritimus flavus, Crithmum chrysanthemum dictus, Raii Hist. 268.  
 157 Brunella Caroliniana, magno flore, dilutè cœruleo; internodiis prælongis.  
 158 — Novæ Angliæ, major; foliis longiùs mucronatis.  
 159 Caulalis Hispanica, Cam. Hort. 37. Fig. xi.  
 160 Corchorus Americana, prælongis foliis, capsula striata, subrotunda, brevi, Pluk. Phyt. Tab 255. f. 4.  
 161 Corona Solis Caroliniana, parvis floribus, folio trinervi amplo, aspero, pediculo alato.  
 162 — — Rapunculi radice, T. Inst. 490  
 163 Dictamnus Creticus, C.B. 222.  
 164 Dorycnium Monspeliensium, Lob. Icon. 2. 51.

- 165 *Elichrysum sylvestre*, latifolium, capitulis conglobatis, C.B. 264.
- 166 *Eupatorium Senecionis facie*, folio *Lamii*, Par. Bat. Ic. 161.
- 167 *Fænum Græcum*, quod *Melilotus corniculis reflexis major*, C.B. 331.
- 168 *Fumaria minor*, tenuifolia, cauliculis procumbentibus & caducis, C.B. 143.
- 169 *Genista Hispanica humilis*, floribus congestis villosis. *Sparto primo affinis*, C.B. 396.
- 170 *Geranium Cicutæ folio*, erectum, Romanum, Bocc. Mus. p. 2, 93.
- 171 — *Columbinum erectum*, *Carolinianum*, floribus purpurascens.
- 172 *Hermannia frutescens*, foliis oblongis, ad apices solummodò ferratis.
- 173 *Jacobæa Africana*, frutescens, flore amplo purpureo elegantissimo, *Senecionis folio*, Volk. Flor. Norimb.
- 174 *Jacobæa Sicula*, *Chrysanthemi facie*, Bocc. rar. p. 66.
- 175 *Lychnidea Virginiana*, *Holosteï ampliore folio*, floribus umbellatis, purpureis.
- 176 *Marrubium album*, foliis amplis, ferè circinatis.
- 177 *Myosotis Hispanica segetum*, T. Inst. 245.
- 178 — *Orientalis perfoliata*, *Lychnidis folio*, T. Cor. 18.
- 179 *Pentaphylloides erectum*, J.B. 2. 398.
- 180 — *Canadense*, folio *Agrimoniæ*. *Sarracen. D. Vaillant.*
- 181 — *Fragariæ folio*, Bot. Monsp. App. 309.
- 182 — — — & crenis obtusioribus.
- 183 *Quinquefolium rectum luteum*, C.B. 325.
- 184 — quod *Pentaphyllum*, seu potius *Heptaphyllum rectum*, caule rubro, hirsutius, H. Cath. Boerh. Ind. 2 40.

- 185 — minus, flore pallidè luteo, T. Inst. 297.  
 186 — folio argenteo, C.B. 325.  
 187 *Phlomis fruticosa*, humilis, latifolia, candidissima ;  
 floribus luteis.  
 188 — *Orientalis*, lutea, herbacea, latifolia, verticil-  
 lata.  
 189 *Sclarea rugoso*, verrucoso & laciniato folio, T. Inst.  
 180.  
 190 *Senecio Ægyptius*, folio *Matricariæ*, Boerh. Ind.  
 1. p. 40.  
 191 *Sideritis arvensis*, latifolia, glabra, C.B. 233.  
 192 — *Vulgaris hirsuta erecta*, C.B. Ib.  
 193 *Serratula annua*, feminibus nitidissimis ad basin  
 compressis. *Senecio carduus Apulus*, Col. p. 1,  
 32.  
 194 — —, — —, majoribus & minimè compressis.  
*Crupina Belgarum* vulgò.  
 195 *Teucrium Bæticum*, Clus. Hisp. 229.  
 196 — —, humilior & ramosior, folio subrotundo.  
 197 *Thlaspi arvense*, *Vaccariæ incano* folio, majus,  
 C.B. 206.  
 198 —, *Vaccariæ incano* folio, minus, C.B. Ib.  
 199 *Virga aurea*, *Canadensis*, *Asterisci* folio, Par. Bat. 244.  
 200 — —, montana, *Scrophulariæ* folio, Icon. Robert.  
 Pluk. Phyt. Tab. 235. f. 3.